

Movements of foraging king penguins through marine mesoscale eddies

Cédric Cotté^{1,*}, Young-Hyang Park², Christophe Guinet¹
and Charles-André Bost¹

¹Centre d'Études Biologiques de Chizé, Centre National de la Recherche Scientifique,
79360 Villiers en Bois, France

²LOCEAN/USM402, Département Milieux et Peuplements Aquatiques, Muséum National d'Histoire Naturelle,
43 rue Cuvier, 75005 Paris, France

Despite increasing evidence that marine predators associate with mesoscale eddies, how these marine features influence foraging movements is still unclear. This study investigates the relationship of at-sea movements of king penguins to mesoscale eddies using oceanographic remote sensing and movement data from 43 individual trips over 4 years. Simultaneous satellite measurements provided information on gradients of sea surface temperature and currents associated with eddies determined from altimetry. Penguins tended to swim rapidly with currents as they travelled towards foraging zones. Swimming speed indicative of foraging occurred within mesoscale fronts and strong currents associated with eddies at the Polar Front. These results demonstrate the importance of mesoscale eddies in directing foraging efforts to allow predators to rapidly get to rich areas where high concentrations of prey are likely to be encountered. When returning to the colony to relieve the incubating partner or to feed the chick, the birds followed a direct and rapid path, seemingly ignoring currents.

Keywords: king penguin; foraging; mesoscale eddies; marine currents; fronts; oceanographic features

1. INTRODUCTION

The open ocean is a heterogeneous environment that is characterized by a variety of physical features. The complexity of this environment results in patchy production over a very broad range of scales in both space and time. Pelagic predators such as seabirds and turtles are reported to actively use particular oceanic features to travel (Weimerskirch *et al.* 2000; Luschi *et al.* 2003; Gaspar *et al.* 2006) and search efficiently for their food supply (Nevitt *et al.* 1995). Owing to the patchiness in marine resources, these mechanisms are expected to have a crucial importance on the foraging strategy of marine predators.

Although they range widely over the open ocean, predators are thought to concentrate their foraging activity according to mesoscale features such as eddies where prey biomass is likely to be elevated (Nel *et al.* 2001; Polovina *et al.* 2006; Ream *et al.* 2005). Mesoscale eddies are suspected to be of primary importance for pelagic ecosystems because they serve as centres of biological production (Strass *et al.* 2002). Recent work suggests the importance of eddies in providing foraging opportunities for flying birds (Nel *et al.* 2001; Weimerskirch *et al.* 2004) as well as for other predators such as sea turtles (Ferraro *et al.* 2004) although other uses have been reported that are life-stage dependent (Hays *et al.* 2006).

To date, no study has investigated how penguin species use this dynamic environment to travel and search for profitable patches. These diving seabirds face different environmental constraints than dynamic soaring seabirds, in that they have to deal with the high cost of swimming

when compared with the low cost of soaring and gliding (Costa 1991). As a result, their foraging range is restricted during the breeding season (Wilson 1995). King penguins (*Aptenodytes patagonicus*) are one of the major avian predators of the Southern Ocean in terms of population size and prey consumption (Woehler 1995). During the breeding season, these diving birds mainly exploit the Polar Front, where they feed on pelagic fishes (Bost *et al.* 1997; Guinet *et al.* 1997; Charassin & Bost 2001; Sokolov *et al.* 2006). The Polar Front is a productive circumpolar boundary that corresponds to the northernmost limit of the Antarctic waters (Park *et al.* 1993). As it is also a dynamic front that is characterized by the presence of many mesoscale eddies (Park *et al.* 2002; Kostianoy *et al.* 2004), the behaviour of king penguins offers an ideal opportunity to explore how a swimming predator uses mesoscale eddies to forage.

Firstly, our aim was to assess whether king penguins forage preferentially in mesoscale frontal zones defined by sea surface temperature (SST) gradients, as has been shown for other long-range marine predators (Sims & Quayle 1998). Secondly, we wanted to investigate how movement en route to the Polar Front oriented to current. Here, we examined king penguin movement patterns by investigating swimming speed and direction relative to currents associated with eddies. We predicted that penguins would swim in the same direction as current en route to foraging spots as a mechanism to reduce travel cost.

2. MATERIAL AND METHODS

(a) Datasets

King penguins nesting on Possession Island, Crozet Archipelago were fitted with platform transmitter terminals

* Author for correspondence (cecotte@cebc.cnrs.fr).

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2007.0775> or via <http://www.journals.royalsoc.ac.uk>.

(PTTs; model Kiwisat 101, Sirtrack Ltd., Havelock North, New Zealand) each austral summer (December to March) from 2002 to 2005. A total of 43 breeding king penguins were tracked. The transmission interval was 45 s with an on/off interval of 6 h. The on/off interval was programmed to ensure reception of uplinks during the time of the day that has previously been determined as corresponding to high penguin at-sea activity (Charrassin *et al.* 2002). To ensure that breeding was not disturbed, birds were equipped before departure to sea just after they had been relieved by their partner at the nest. The PTTs weighed 208 g, i.e. approximately 1.8% of body weight, and were hydrodynamically streamlined. They were fitted to the lower back to reduce drag, with the antennae positioned posteriorly (Bannasch *et al.* 1994). The devices were fixed to the back feathers with cyanoacrylate glue (Loctite 401) and secured with cable ties. Antenna-induced effect of satellite transmitters may have deleterious effects on the hydrodynamics of swimming (Wilson *et al.* 2004). In our study, the trip duration of brooding penguins instrumented with PTT was increased only by 9%. The meal size of penguins fitted with PTTs was not different from those of control birds (C. A. Bost 2005, personal observation). Likewise, no differences in breeding failure were reported among control and instrumented birds. Location data were analysed using ELSA software (Argos CLS, Toulouse, France) and custom-written software CROZARG. The accuracy of Argos locations depends on the number of uplinks received. The daily frequency of retained uplinks varied between 8 and 15. Therefore, almost all categories of Argos locations (A, B, 0, 1, 2 and 3, but not Z) were included in the analyses. Swimming speed was conservatively estimated by assuming a straight-line direction and at constant speed between two successive locations (Weavers 1992). As speed errors could come from Argos location inaccuracies (Hays *et al.* 2001), location errors were filtered such that locations indicating a travelling speed higher than 14 km h^{-1} were rejected. This value corresponds to the greatest swimming speed recorded in a king penguin (Kooyman & Davis 1987).

Simultaneous oceanographic data were obtained from satellite measurements. We used 9 km resolution satellite-derived SST from advanced very high resolution radiometer and moderate-resolution imaging spectroradiometer. Although both SST data are available as a daily product, the radiometer does not penetrate cloud cover. Therefore, we used weekly averages as clouds diminished. This temporal resolution was sufficient for studying mesoscale dynamics. To identify mesoscale fronts, we applied a moving window across the grid of SST values. This window permitted slope calculations within a 5×5 neighbourhood calculated by

$$\text{percent slope} = \sqrt{\left(\frac{\delta T}{\delta x}\right)^2 + \left(\frac{\delta T}{\delta y}\right)^2} \times 100, \quad (2.1)$$

where x , y and T indicate longitude; latitude; and SST values, respectively. A low-pass filter (Gaussian method on a 3×3 moving window) was then applied to the new grid of slope values to remove the high-frequency noise and thus smoothed the frontal structures. We considered the surface signature of a Polar Front to be between the 4 and 5°C isotherms (Park *et al.* 1993). Associated with this surface approximation of the Polar Front, the SST gradients reliably reveal the presence of mesoscale fronts (Kostianoy *et al.* 2004).

Weekly maps of gridded sea level anomalies (SLA) at 0.3° resolution on a Mercator projection were used for the periods

of penguin tracking. These maps were made by merging datasets from T/P, ERS-1/2, Jason and Envisat satellite altimeters (Ducet *et al.* 2000). The SLA data were distributed by the archiving, validation and interpretation of satellite oceanographic project of the collecte, localization, satellite (CLS/Centre National d'Études Spatiales, France) for the SSALTO program. Despite possibilities of error due to differences in ground tracks and orbit repeat periods among the several altimeter instruments and to interpolation in the grid construction, merged data have the advantage of resolving mesoscale oceanic processes beyond the capability of a single instrument (Fu *et al.* 2003). Geostrophic currents (0.3° resolution) were computed from SLA as follows:

$$u = -(g/f) \frac{\delta H}{\delta y} \quad \text{and} \quad (2.2)$$

$$v = (g/f) \frac{\delta H}{\delta x}, \quad (2.3)$$

where $g = 980 \text{ cm s}^{-2}$; $f = 2\Omega \sin \Phi$; $\Omega = 7.29 \times 10^{-5} \text{ radians s}^{-1}$; Φ is the latitude; and H is sea level anomaly

(b) Analyses

For each penguin trip, we identified three phases using running averages over five consecutive swimming speeds and compared these running averages with the mean swimming speed for the whole trip. In doing so, we assumed that king penguins would decrease their swimming speed to actively forage, i.e. they would decrease their horizontal (displacements) activity while they increased their vertical (diving) activity (in accordance with Bost *et al.* 1997; Pütz *et al.* 1999; Charrassin *et al.* 2002). These three phases were defined as the following. (i) The transit phase, indicating the travel between the colony and the central phase, where the running averages of swimming speeds were higher than the mean speed of the whole trip. (ii) The central phase, which began at the first episode of slowing down, where the running averages were below the mean speed. Slowing-down episodes in the central phase were frequently separated by speed-up episodes, corresponding to running averages above the overall mean speed. (iii) The return phase, where running averages of swimming speed were higher than the overall mean speed, followed the last slowing-down episode of the central phase. Penguins in incubating and chick-rearing stages were pooled for this analysis. Since slowing-down episodes of the central phase represented intense foraging activity (Bost *et al.* 1997), they were compared statistically with episodes of speeding up in the central phase and also with transit and return phases. To tie these behaviours in with oceanographic data, corresponding analyses were performed using oceanographic parameters (SST, SST gradient and currents from SLA) from measurements taken closest to the penguin's at-sea locations. Bearing deviations between penguin and underlying current directions, varying from 0° to 360° , were also investigated. A bearing deviation from 0° to 90° and from 270° to 360° signified similar directions for a penguin and the current (positive rheotaxis), while a bearing deviation from 90° to 270° corresponded to opposing directions (negative rheotaxis; Ream *et al.* 2005). Similar (0° – 90° and 270° – 360°) directions suggested that currents assisted the directional movements of the penguins whereas opposing (90° – 270°) directions suggested opposing forces. Net king penguin speed and movement are defined by parameters \bar{u}_N (east–west) and

Table 1. Sea surface temperature gradients (gSST, means \pm s.e.) for all locations of the 43 king penguin trips.

trip phases	gSST (%)	U-test (versus slowing-down episodes)
transit	2.13 ± 1.28	$p < 0.001$
central		
slowing-down episodes	2.95 ± 1.80	
speed-up episodes	2.65 ± 1.61	$p = 0.023$
return	2.19 ± 1.36	$p < 0.001$

\bar{v}_N (north–south), and currents by \bar{u}_C (east–west) and \bar{v}_C (north–south), with the total vectors estimated as $\sqrt{(\bar{u}^2 + \bar{v}^2)}$. Net penguin speed was the resultant of swimming and underlying current speeds. Thus, we calculated the real swimming speed vector (in terms of amplitude and direction) of penguins by $\bar{u}_R = \bar{u}_N - \bar{u}_C$ and $\bar{v}_R = \bar{v}_N - \bar{v}_C$. We used circular statistics to estimate mean length (ML) of mean swimming and current vectors, and also direction confidence interval (CI) from bootstrap method for each phase.

3. RESULTS

(a) King penguin trips and mesoscale fronts

For all trips, penguins tended to slow down when they encountered the steepest SST gradient (table 1). Steep SST gradients are indicators of mesoscale fronts, where physical and biological processes occur. This relationship suggests that penguins were slowing down to forage at these mesoscale fronts while the rest of the trip, i.e. transit and return phases and also speed-up episodes during the central phase, tended to be associated with shallower SST gradients. These successive phases are detailed for one typical trip (figure 1) where the several slowing-down episodes were clearly associated with steeper SST gradients.

(b) King penguin movements and mesoscale currents

Frequencies of estimated mesoscale current directions in the whole area, defined by tracking bird ranges, show a slight northeastward direction during the transit and return phases (between Crozet Island and the Polar Front), and a slight westward direction during the central phase (within the Polar Front; figure 2). The mesoscale current situation is highly changing due to eddies, variable in both time and amplitude. Currents encountered along the tracks were significantly stronger during slowing-down episodes than during the rest of trip (table 2). Bearing deviations during the transit phase indicate that birds tended to swim in a similar direction as the current (bird direction, CI = 170° – 178° , ML = 0.52; current direction, CI = 174° – 185° , ML = 0.33; 0° and 180° being north and south direction, respectively; Watson test $U^2 = 0.21$, $p > 0.1$; table 3 and figure 3). This trend disappeared during slowing-down episodes characteristic of the central phase, where birds had no preferred direction (bird direction, CI = 169° – 214° , ML = 0.09), and suggesting that birds no longer swam with the current (CI = 128° – 150° , ML = 0.16). Speed-up episodes of the central phase, where penguins left a foraging place to another one, indicate




Figure 1. Parameters for a king penguin track with (a) penguin swimming speed from running average over five positions (horizontal line indicates the mean speed for the whole trip), (b) sea surface temperature, SST (horizontal line indicates northern limit of the Polar Front) and (c) sea surface temperature gradients (gSST). Grey bars indicate slowing-down episodes during the central phase, defined as swimming speeds below the mean. Slowing-down episodes correspond to bouts of intense foraging (Bost et al. 1997).

that birds tended to swim with currents as in the transit phase (bird direction, CI = 170° – 189° , ML = 0.4; current direction, CI = 157° – 199° , ML = 0.17; $U^2 = 0.07$, $p > 0.1$). During the return phase, penguins tended to swim against the current (bird direction, CI = 345° – 353° , ML = 0.51; current direction, CI = 144° – 172° , ML = 0.14; $U^2 = 0.47$, $p < 0.001$). Currents speeds were at least one order of magnitude lower than penguin swimming speeds, suggesting that they could easily swim against the current (table 2). Swimming speeds during the transit phase and speed-up episodes of the central phase were similar, while they were faster on average during the return phase.

Figure 4 illustrates penguin movements in relation to currents. These journeys of three different seabirds corresponded to a situation where an eddy remained static for at least one month. The birds exhibited similar trip patterns in response to a similar mesoscale pattern of marine circulation: southward transits and northward returns were in straight lines between the colony and an eddy while once at the edge of this anticyclonic eddy the three trips were clockwise, i.e. in a similar direction as the prevailing current.




Figure 2. Frequencies of directions of marine currents, estimated from satellite altimetry, in the areas prospected by king penguins (a) during transit and return phases and (b) during the central phase. Limits of the areas are defined by tracking bird ranges.

4. DISCUSSION AND CONCLUSIONS

There is increasing evidence that a wide variety of marine vertebrates aggregate in association with mesoscale eddies (e.g. Nel *et al.* 2001). However, very few studies go further than documenting an empirical association. Because eddies are characterized by strong gradients in water mass properties, especially SST gradients in the Polar Front (Kostianoy *et al.* 2004), it is relevant to assess quantitative associations between penguin activity (travelling and foraging) and such gradients (SST gradients and currents). Our analysis suggests that penguin swimming behaviour relative to the current changes at mesoscale fronts, where penguins are known to increase their diving activity and food intake (Bost *et al.* 1997; Charassin & Bost 2001). The changes were especially marked at the periphery of eddies, suggesting that the foraging activity occurred more at the edge of eddies than the centre (Nel *et al.* 2001; Weimerskirch *et al.* 2004; Polovina *et al.* 2006). Besides the broad strategy of king penguins to go to the Polar Front, these results confirm the importance of considering the influence of finer-scale physical parameters. In particular, frontal meanders of the Polar Front created by eddies give rise to mesoscale convergence and divergence. These features can concentrate zooplankton and micronekton (Pakhomov & Fronemann 2000), preyed upon by fish such as myctophids, which opportunistically forage on mesozooplankton (Pakhomov *et al.* 1996) and presumably attract king penguins as well as others upper-trophic predators (Hyrenbach *et al.* 2006). Eddy structures are beneficial to predators. Upwelling occurs at the centre, whereas convergence (horizontal concentration of productivity and preys) occurs at the edge of cyclonic (cold-core) eddy. The edge of anticyclonic (warm-core) eddy presents an elevation of water masses, leading to a more accessible concentration of prey at shallow depths (Polovina *et al.* 2006). Several foraging predators including penguins have been shown to forage in association with thermoclines, strong physical discontinuities that concentrate potential prey items (Charassin & Bost 2001). Were thermoclines to occur at shallower depths, this would be beneficial to penguins since shallower dives would be required for prey capture. Despite their ephemeral nature, these mesoscale fronts may provide useful feeding grounds for top predators, particularly in pelagic systems where prey patches are overdispersed (Nel *et al.* 2001; Weimerskirch 2007). Furthermore, these

mesoscale features are relatively common within the Polar Front and may be particularly important for predators such as king penguins, which are not able to cover vast distance during the constraints of the breeding season. Although transitory eddies and associated ephemeral fronts dominate the Polar Front, some marked static physical features can nevertheless occur. Since birds tend to travel to similar locations to forage, even over vast distances, persisting eddies may be ecologically relevant. These longer term productive features are likely to constitute favourable and above all predictable feeding grounds for highly constrained predators like penguins. Persisting eddies can have their own production and food chain development in contrast to usual transitory features that simply aggregate planktonic organisms (Palacios *et al.* 2006).

Recently, studies on marine predator movements have particularly emphasized the mechanical influence of currents with respect to their deflecting action, forcing animals to drift away from the most suitable route (Gaspar *et al.* 2006; Girard *et al.* 2006). Turtles were thus reported to be 'caught' by the circulation, concentrating their activity on swimming vertically rather than horizontally to forage (Luschi *et al.* 2003). While these findings concerned predators with a swimming speed of the same order of magnitude as the current speed, our results are focusing on a situation where penguin movements are substantially faster than the ambient currents (by a factor of 10 at least). In this situation, the predator movements are indeed less impacted by the deflecting action of currents. This important difference was due to the following reasons. Firstly, their trips under breeding constraints were made with a sustained swimming speed comparable with breeding Antarctic fur seals (1–8 km h⁻¹ from Boyd *et al.* 1995) and higher than most long-range foraging predators such as turtles (0.7–3.5 km h⁻¹ from Luschi *et al.* 1998; Polovina *et al.* 2004; Ferraroli *et al.* 2004, 2006; Girard *et al.* 2006), tuna (approx. 1.5 km h⁻¹ from Block *et al.* 1998) and non-breeding seals (0.3–2.2 km h⁻¹ from Ream *et al.* 2005; Austin *et al.* 2006). Secondly, the currents in the Polar Frontal zone are weaker than that in northern fronts that constitute the major portion of the antarctic circumpolar current (Park *et al.* 2002) and also weaker than that in most regions where animal movements have been studied (Luschi *et al.* 2003; Ferraroli *et al.* 2004; Polovina *et al.* 2006).




Figure 3. Frequencies of bearing deviations between penguin and underlying current direction for all locations of the 43 king penguin trips during (a) transit phase, (b) speed-up and (c) slowing-down episodes of the central phase and (d) return phase. Arrows indicate the referential swimming direction of penguins and the dotted lines separate similar (270° – 90°) and opposing (90° – 270°) directions.

Table 2. Means \pm s.e. of penguin swimming and underlying current speeds for all locations of the 43 king penguin trips.

trip phases	swimming speed (km h^{-1})	<i>U</i> -test (versus slowing- down episodes)	current speed (km h^{-1})	<i>U</i> -test (versus slowing- down episodes)
transit	5.8 ± 4.0	$p < 0.001$	0.23 ± 0.14	$p < 0.001$
central				
slowing-down episodes	3.9 ± 3.6		0.31 ± 0.18	
speed-up episodes	5.7 ± 4.1	$p < 0.001$	0.23 ± 0.13	$p < 0.001$
return	6.4 ± 4.8	$p < 0.001$	0.24 ± 0.14	$p < 0.001$

Despite penguin swimming speed substantially greater than current speed, our study shows that king penguins shifted their movement behaviour in relation to currents through the distinct phases of their trips. Importantly, penguin movements appeared to be related to currents as they commute from the colony to foraging places, but are unrelated during actual foraging. Our results support the hypothesis, also suggested for turtles, that penguins may take advantage of current flow to increase their speed without increasing energy cost (Hays *et al.* 1999). Breeding penguins leave the colony after several weeks of fasting and they have to rapidly rebuild their own energetic reserves before stocking food for their offspring during chick rearing (Kooyman *et al.* 1992). Once penguins slowed down to forage, swimming direction no longer correlates with current direction, despite stronger currents at the periphery of eddies. This behavioural shift is probably due to the

Table 3. Trip proportion presenting similar penguin–current directions (i.e. bearing deviation between 270° and 90°) from the mean bearing deviation per phase for each of the 43 king penguin trips.

trip phases	penguin–current similar directions (%)
transit	74
central	
slowing-down episodes	53
speed-up episodes	63
return	39

increase in foraging effort (Bost *et al.* 1997; Charassin & Bost 2001), and also due to the optimal exploitation of the mesoscale fronts until the benefit from staying in this area either drops below some threshold or they replenish their




Figure 4. SLA and geostrophic currents with successive tracks from three different penguins (a) first, 28 January 2004 to 10 February 2004 (b) second, 2 February 2004 to 24 February 2004 and (c) third, 11 February 2004 to 25 February 2004. The star indicates the position of Crozet Archipelago. Arrows give trip directions.

reserves. During the return phase, where diving rate and food intake decrease drastically (Bost *et al.* 1997; Charrassin *et al.* 2002), penguins came back rapidly to the colony owing to the breeding constraints, i.e. partner relieve on egg or chick, or chick feeding, effectively swimming up stream to do so. Since they used mainly southward currents while travelling towards the Polar Front, the penguins had to face these currents when they came back, which explain the higher proportion of opposing currents encountered during the return.

Mesoscale eddies are thus suggested to be particularly important for king penguins through (i) the creation of mesoscale fronts that are potential feeding grounds in the vicinity of the Polar Front and (ii) providing king penguins with either a faster means of reaching their goal or a directional cue to get there. By contrast to relatively slow marine turtles, affected by the drifting action of strong currents (Gaspar *et al.* 2006; Girard *et al.* 2006), faster penguins highly constrained in space and time may take advantage of these currents as an aid to progress rapidly to favourable fronts associated with the mesoscale eddy activity. Despite certain technical limitations on the availability of physical measurements from the surface (e.g. cloudiness), remote sensing provides undoubtedly valuable information on the relationship between at-sea animal activity and its environment. Furthermore, mechanistic studies are still required to better understand the causal biophysical interaction and the link between lower and upper trophic system constituents that take place at these mesoscale eddies (Hyrenbach *et al.* 2006). The present study provides new insights into the influence of eddy activity on the behaviour of marine predators, particularly in high latitudes. It highlights the role that mesoscale features play in marine ecosystems as well as the lack of information on mechanisms driving the biological production and prey aggregation associated with these features. Further studies will examine diving and feeding activity associated with eddies during king penguin foraging trips. In this mesoscale context, the role of foraging experience through tracking successive trips of the same birds should be especially promising.

This study was approved by the ethics committee of the French Polar Institute (IPEV). We thank P.-M. Theveny for helping with data analysis and L. G. Hasley, H. Weimerkirch,

J.-B. Charrassin, Y. Cherel and D. Pinaud for their helpful comments on an earlier version of the manuscript. Special thanks to A. Pape for his help with the English. This study was carried out in the framework of GICC and REMIGE (ANR 2005 Biodiv-11) programs. The IPEV and the TAAF provided financial and logistical support. We thank Gabrielle Nevitt and two anonymous referees for their suggestive and helpful comments that substantially improve the manuscript.

REFERENCES

- Austin, D., Bowen, W. D., McMillan, J. I. & Iverson, S. J. 2006 Linking movement, diving, and habitat to foraging success in a large marine predator. *Ecology* **87**, 3095–3108. (doi:10.1890/0012-9658(2006)87[3095:LMDAHT]2.0.CO;2)
- Bannasch, R., Wilson, R. P. & Culik, B. 1994 Hydrodynamic aspects of design and attachment of a back-mounted device in penguins. *J. Exp. Biol.* **194**, 83–96.
- Block, B. A., Dewar, H., Farwell, C. & Prince, E. D. 1998 A new satellite technology for tracking the movements of Atlantic bluefin tuna. *Proc. Natl Acad. Sci. USA* **95**, 9384–9389. (doi:10.1073/pnas.95.16.9384)
- Bost, C. A. *et al.* 1997 Foraging habitat and food intake of satellite-tracked king penguins during the austral summer at Crozet Archipelago. *Mar. Ecol. Prog. Ser.* **150**, 21–33.
- Boyd, I. L., Reid, K. & Bevan, R. M. 1995 Swimming speed and allocation of time during the dive cycle in Antarctic fur seals. *Anim. Behav.* **50**, 769–784. (doi:10.1016/0003-3472(95)80137-5)
- Charrassin, J. B. & Bost, C. A. 2001 Utilization of the oceanic habitat by king penguins over the annual cycle. *Mar. Ecol. Prog. Ser.* **221**, 285–297.
- Charrassin, J. B., Le Maho, Y. & Bost, C. A. 2002 Seasonal changes in the diving parameters of king penguins (*Aptenodytes patagonicus*). *Mar. Biol.* **141**, 581–589. (doi:10.1007/s00227-002-0843-4)
- Costa, D. P. 1991 Reproductive and foraging energetics of high latitude penguins, albatrosses and pinnipeds: implications for life history patterns. *Am. Zool.* **31**, 11–130.
- Ducet, N., Le Traon, P. Y. & Reverdin, G. 2000 Global high resolution mapping of ocean circulation from TOPEX/Poseidon and ERS-1/2. *J. Geophys. Res.* **105**, 19 477–19 498. (doi:10.1029/2000JC900063)
- Ferraroli, S., Georges, J. Y., Gaspar, P. & Le Maho, Y. 2004 Endangered species: where leatherback turtles meet fisheries. *Nature* **429**, 521–522. (doi:10.1038/429521a)
- Fu, L.-L., Stammer, D., Leben, R. R. & Chelton, D. B. 2003 Improved spatial resolution of ocean surface topography

- from the T/P-Jason-1 altimeter mission. *EOS Trans. Am. Geophys. Union* **84**, 241. (doi:10.1029/2003EO260002)
- Gaspar, P., Georges, J. Y., Fossette, S., Lenoble, A., Ferraroli, S. & Le Maho, Y. 2006 Marine animal behaviour: neglecting ocean currents can lead us up the wrong track. *Proc. R. Soc. B* **273**, 2697–2702. (doi:10.1098/rspb.2006.3623)
- Girard, C., Sudre, J., Benhamou, S., Roos, D. & Luschi, P. 2006 Homing in green turtles *Chelonia mydas*: oceanic currents act as a constraint rather than as an information source. *Mar. Ecol. Prog. Ser.* **322**, 281–289.
- Guinet, C., Koudil, M., Bost, C. A., Durbec, J. P., Georges, J. Y., Mouchot, M. C. & Jouventin, P. 1997 Foraging behaviour of satellite-tracked king penguins in relation to sea-surface temperatures obtained by satellite telemetry at Crozet Archipelago, a study during three austral summers. *Mar. Ecol. Prog. Ser.* **150**, 11–20.
- Hays, G. C., Luschi, P., Papi, F., del Seppia, C. & Marsh, R. 1999 Changes in behaviour during the interesting period and postnesting migration for Ascension Island green turtles. *Mar. Ecol. Prog. Ser.* **180**, 263–273.
- Hays, G. C., Åkesson, S., Godley, B. J., Luschi, P. & Santidrian, P. 2001 The implications of location accuracy for the interpretation of satellite tracking data. *Anim. Behav.* **61**, 1035–1040. (doi:10.1006/anbe.2001.1685)
- Hays, G. C., Hobson, V. J., Metcalfe, J. D., Righton, D. & Sims, D. W. 2006 Flexible foraging movements of leatherback turtles across the North Atlantic Ocean. *Ecology* **87**, 2647–2656. (doi:10.1890/0012-9658(2006)87[2647:FFMOLT]2.0.CO;2)
- Hyrenbach, K. D., Veit, R. R., Weimerskirch, H. & Hunt Jr, G. L. 2006 Seabirds associations with mesoscale eddies: the subtropical Indian Ocean. *Mar. Ecol. Prog. Ser.* **324**, 271–279.
- Kooyman, G. L. & Davis, R. W. 1987 Diving behavior and performance, with special reference to penguins. In *Seabirds feeding ecology and role in marine ecosystems* (ed. J. P. Croxall), pp. 63–75. Cambridge, UK: Cambridge University Press.
- Kooyman, G. L., Cherel, Y., Le Maho, Y., Croxall, J. P., Thorson, P. H., Ridoux, V. & Kooyman, C. A. 1992 Diving behaviour and energetics during foraging cycles in king penguins. *Ecol. Monogr.* **62**, 143–163. (doi:10.2307/2937173)
- Kostianoy, A. G., Ginzburg, A. I., Frankignoulle, M. & Dellile, B. 2004 Fronts in the Southern Indian Ocean as inferred from satellite sea surface temperature data. *J. Mar. Syst.* **45**, 55–73. (doi:10.1016/j.jmarsys.2003.09.004)
- Luschi, P., Hays, G. C., del Seppia, C., Marsh, R. & Papi, F. 1998 The navigational feats of green turtles migrating from Ascension Island investigated by satellite telemetry. *Proc. R. Soc. B* **265**, 2279–2284. (doi:10.1098/rspb.1998.0571)
- Luschi, P., Sale, A., Mencacci, R., Hughes, G. R., Lutjeharms, J. R. E. & Papi, F. 2003 Current transport of leatherback sea turtles (*Dermochelys coriacea*) in the ocean. *Proc. R. Soc. B* **270**(Suppl.), 129–132. (doi:10.1098/rsbl.2003.0036)
- Nel, D. C., Lutjeharms, J. R. E., Pakhomov, E. A., Anson, I. J., Ryan, P. G. & Klages, N. T. W. 2001 Exploitation of mesoscale oceanographic features by grey-headed albatross *Thalassarche chrysostoma* in the Southern Indian Ocean. *Mar. Ecol. Prog. Ser.* **217**, 15–26.
- Nevitt, G. A., Veit, R. R. & Kareiva, P. 1995 Dimethyl sulphide as a foraging cue for Antarctic Procellariiform seabirds. *Nature* **376**, 680–682. (doi:10.1038/376680ao)
- Pakhomov, E. A. & Fronemann, P. W. 2000 Composition and spatial variability of macrozooplankton and micronekton within the Antarctic Polar Frontal Zone of the Indian Ocean during austral summer 1997. *Polar Biol.* **23**, 410–419. (doi:10.1007/s003000050462)
- Pakhomov, E. A., Perissinotto, R. & McQuaid, C. D. 1996 Prey composition and daily rations of myctophid fishes in the Southern Ocean. *Mar. Ecol. Prog. Ser.* **134**, 1–14.
- Palacios, D. M., Bograd, S. J., Foley, D. G. & Schwing, F. B. 2006 Oceanographic characteristics of biological hot spots in the North Pacific: a remote sensing perspective. *Deep Sea Res. II* **53**, 250–269. (doi:10.1016/j.dsr2.2006.03.004)
- Park, Y. H., Gambérini, L. & Charriaud, E. 1993 Frontal structure, water masses, and circulation in the Crozet Basin. *J. Geophys. Res.* **98**, 12 361–12 385.
- Park, Y. H., Pollard, R. T., Read, J. F. & Lebourcier, V. 2002 A quasi-synoptic view of the frontal circulation in the Crozet Basin during the Antares-4 cruise. *Deep Sea Res. II* **49**, 1823–1842. (doi:10.1016/S0967-0645(02)00014-0)
- Polovina, J. J., Balazs, G. H., Howell, E. A., Parker, D. M., Seki, M. P. & Dutton, M. P. 2004 Forage and migration habitat of loggerhead (*Caretta caretta*) and olive ridley (*Lepidochelys olivacea*) sea turtles in the central North Pacific Ocean. *Fish. Oceanogr.* **13**, 36–51. (doi:10.1046/j.1365-2419.2003.00270.x)
- Polovina, J. J., Uchida, I., Balazs, G. H., Howell, E. A., Parker, D. M. & Dutton, M. P. 2006 The Kuroshio extension bifurcation region: a pelagic hotspot for juvenile loggerhead sea turtles. *Deep Sea Res. II* **53**, 326–339. (doi:10.1016/j.dsr2.2006.01.006)
- Pütz, K., Ropert-Coudert, Y., Charrassin, J. B. & Wilson, R. P. 1999 Foraging areas of King penguins *Aptenodytes patagonicus* breeding at Possession Island, southern Indian Ocean. *Mar. Ornithol.* **27**, 77–84.
- Ream, R. R., Sterling, J. T. & Loughlin, T. R. 2005 Oceanographic features related to northern fur seal migratory movements. *Deep Sea Res. II* **52**, 823–843. (doi:10.1016/j.dsr2.2004.12.021)
- Sims, D. W. & Quayle, V. A. 1998 Selective foraging behaviour of basking sharks on zooplankton in a small-scale front. *Nature* **393**, 460–464. (doi:10.1038/30959)
- Sokolov, S., Rintoul, S. R. & Wienecke, B. 2006 Tracking the Polar Front south of New Zealand using penguin dive data. *Deep Sea Res. I* **53**, 591–607. (doi:10.1016/j.dsr.2005.12.012)
- Strass, V. H., Naveira Garabato, A. C., Pollard, R. T., Fischer, H. I., Hense, I., Allen, J. T., Read, J. F., Leach, H. & Smetacek, V. 2002 Mesoscale frontal dynamics: shaping the environment of primary production in the Antarctic Circumpolar Current. *Deep Sea Res. II* **49**, 3735–3769. (doi:10.1016/S0967-0645(02)00109-1)
- Weavers, B. W. 1992 Seasonal foraging ranges and travels at sea of little blue penguins *Eudyptula minor*, determined by radiotracking. *Emu* **91**, 302–307.
- Weimerskirch, H. 2007 Are seabirds foraging for unpredictable resources? *Deep Sea Res. II* **54**, 211–223. (doi:10.1016/j.dsr.2.2006.11.013)
- Weimerskirch, H., Guionnet, T., Martin, J., Shaffer, S. A. & Costa, D. P. 2000 Fast and fuel efficient? Optimal use of wind by flying albatrosses. *Proc. R. Soc. B* **267**, 1869–1874. (doi:10.1098/rspb.2000.1223)
- Weimerskirch, H., Le Corre, M., Jacquemet, S., Potier, M. & Marsac, D. P. 2004 Foraging strategy of a top predator in tropical waters: great frigatebirds in the Mozambique Channel. *Mar. Ecol. Prog. Ser.* **275**, 297–308.
- Wilson, R. P. 1995 Foraging ecology. In *The penguins* (ed. T. D. Williams), pp. 81–106. Oxford, UK: Oxford University Press.
- Wilson, R. P., Kreye, J. M., Lucke, K. & Urquhart, H. 2004 Antennae on transmitters on penguins: balancing energy budgets on the high wire. *J. Exp. Biol.* **207**, 2649–2662. (doi:10.1242/jeb.01067)
- Woehler, E. J. 1995 Consumption of Southern Ocean resources by penguins. In *The penguins, ecology and management* (eds P. Dann, I. Norman & P. Reilly), pp. 267–291. Chipping Norton, UK: Surrey Beatty and Sons.